

On strong n -perfect rings

Abdellatif Jhilal and Najib Mahdou

Department of Mathematics, Faculty of Science and Technology of Fez,
Box 2202, University S. M. Ben Abdellah Fez, Morocco,
Jhilalabdo@hotmail.com
mahdou@hotmail.com

Abstract.

In this paper we introduce the notion of “strong n -perfect rings” which is in some way a generalization of the notion of “ n -perfect rings”. We are mainly concerned with those class of rings in the context of pullbacks. Also we exhibit a class of n -perfect rings that are not strong n -perfect rings. Finally, we establish the transfer of this notion to the direct product notions.

Key Words. Perfect ring, n -perfect ring, strong n -perfect property, (n, d) -property, pullback ring.

1 Introduction

Throughout this work, all rings are commutative with identity element, and all modules are unitary. Let R be a ring and let M be an R -module. As usual we use $\text{pd}_R(M)$ and $\text{fd}_R(M)$ to denote the usual projective and flat dimensions of M , respectively. $\text{gldim}(R)$ and $\text{wdim}(R)$ are, respectively, the classical global and weak dimension of R . If R is an integral domain, we denote its quotient field by $\text{qf}(R)$.

A ring R is perfect if every flat R -module is projective R -module. The pioneering work on perfect rings was done by Bass [2] and most of the principal characterizations of perfect rings are contained in Theorem P from that paper.

Recently, Enochs, et al. in his work [9] extended the notion of perfect rings to n -perfect rings, such that a ring is called n -perfect, if every flat module has projective dimension less or equal than n [9, Definition 1.1].

The classes of rings we will define here are in some ways generalization of the notion of “ n -perfect rings”. Let n be a positive integer. A commutative ring R is called a strong n -perfect ring if any R -module of flat dimension less or equal than n have projective dimension less or equal than n . Note that if $n = 0$ then the strong 0-perfect rings are the perfect rings. And it is trivial to remark that every strong n -perfect ring is also an n -perfect ring.

In this paper, we investigate the transfer of strong n -perfect property in some know ring constructions. Namely, we study the strong n -perfect property of pullbacks (Theorem 2.3) and of finite direct product (Theorem 2.16). See [10] for a quite general development of properties of pullbacks. And Theorem 2.3 enable us to give a class of strong n -perfect rings (Example 2.6).

We also show that, in a particular cases, the relationship between the strong n -perfect property and (n, d) -property (Theorem 2.7), for background on (n, d) -rings we refer the reader to [4, 5, 7, 15].

As the strong n -perfect property is a general of the n -perfect property, it is important to give examples of n -perfect rings which are not strong n -perfect rings. This is given in Example 2.13.

Finally, the Proposition 2.14 allow us to study the strong n -perfect property of some particular rings.

2 Main results

We begin by the following change of rings result for strong n -perfect property, in which we use the notion of flat epimorphism of rings, which is defined as follows: Let $\Phi : A \rightarrow B$ be a ring homomorphism. B (or Φ) is called a flat epimorphism of A , if B is a flat A -module and Φ is an epimorphism, that is, for any two ring homomorphism $B \begin{smallmatrix} \xrightarrow{f} \\ \xrightarrow{g} \end{smallmatrix} C$, C a ring, satisfying $f \circ \Phi = g \circ \Phi$, we have $f = g$ [12, pages 13-14]. For example $S^{-1}A$ is a flat epimorphism of A for every multiplicative set S of A . Also, the quotient ring A/I is a flat epimorphism of A for every pure ideal I of A , that is, A/I is a flat A -module [12, Theorem 1. 2. 15]

Proposition 2.1 *Let A and B be two rings such that $\Phi : A \rightarrow B$ be a flat epimorphism of A . Then:*

If A is a strong n -perfect ring then B is a strong n -perfect ring.

Proof. : Let M be an B -module such that $\text{fd}_B(M) \leq n$. Our aim is to show that $\text{pd}_B(M) \leq n$.

By hypothesis we have $\text{Tor}_k^A(M, B) = 0$ for all $k > 0$, so for any B -module N , we have from [3, Proposition 4.1.3].

$$(*) \quad \text{Ext}_A^{n+1}(M, N \otimes_A B) \cong \text{Ext}_B^{n+1}(M \otimes_A B, N \otimes_A B)$$

And [12, Theorem 1.2.19] give $M \otimes_A B \cong M$ and $N \otimes_A B \cong N$.

On the other hand $\text{fd}_A(M) \leq \text{fd}_B(M) \leq n$ [3, Exercise 10.p.123]. Thus $\text{pd}_A(M) \leq n$ since A is a strong n -perfect ring. Therefore $\text{Ext}_B^{n+1}(M, N) = 0$ by $(*)$ and so $\text{pd}_B(M) \leq n$. ■

As $S^{-1}A$ and A/I are the particular cases of flat epimorphism of A for any multiplicative set S of A and for any pure ideal I of A . we have the following results.

Corollary 2.2 1. *For every multiplicative set S of a strong n -perfect ring A , $S^{-1}A$ is a strong n -perfect ring ($n \geq 0$).*

2. *If A is a strong n -perfect ring, then so is the quotient ring A/I for every pure ideal I of A ($n \geq 0$).*

Now we give our main result of this paper.

Theorem 2.3 *Let $A \hookrightarrow B$ be an injective flat ring homomorphism and let Q be a pure ideal of A such that $QB = Q$. Then:*

1. *If B is a strong n -perfect ring. Then:
 A/Q is a strong n -perfect ring if and only if A is a strong n -perfect ring.*
2. *If $B = S^{-1}A$, where S is a multiplicative set of A . Then:
 A is a strong n -perfect ring if and only if B and A/Q are a strong n -perfect rings.*

Before proving this theorem, we establish the following lemmas.

Lemma 2.4 [7, Lemma 2.5] *Let $A \hookrightarrow B$ be an injective flat ring homomorphism and let Q be an ideal of A such that $QB = Q$. Let E be an A -module such that $E \otimes_A B$ is B -flat. Then:*

$$\text{pd}_A(E) \leq d \Leftrightarrow \text{pd}_B(E \otimes_A B) \leq d \text{ and } \text{pd}_{A/Q}(E \otimes_A A/Q) \leq d$$

Lemma 2.5 *Let $A \hookrightarrow B$ be an injective flat ring homomorphism and let Q be a pure ideal of A such that $QB = Q$. Let E be an A -module. Then:*

1. $\text{fd}_A(E) \leq d \Leftrightarrow \text{fd}_B(E \otimes_A B) \leq d$ and $\text{fd}_{A/Q}(E \otimes_A A/Q) \leq d$
2. $\text{pd}_A(E) \leq d \Leftrightarrow \text{pd}_B(E \otimes_A B) \leq d$ and $\text{pd}_{A/Q}(E \otimes_A A/Q) \leq d$

Proof. Similar to the proof of Lemma 2.4. ■

Proof of theorem 2.3:

- 1) If A is a strong n -perfect ring, then so is A/Q by Corollary 2.2(2) since Q is a pure ideal of A . Conversely, assume that B and A/Q are a strong n -perfect rings. Let M be an A -module such that $\text{fd}_A(M) \leq n$. Then $\text{fd}_B(M \otimes_A B) \leq n$ and $\text{fd}_{A/Q}(M \otimes_A A/Q) \leq n$ by lemma 2.5.1. Hence, $\text{pd}_B(M \otimes_A B) \leq n$ and $\text{pd}_{A/Q}(M \otimes_A A/Q) \leq n$, since B and A/Q are a strong n -perfect rings. So that $\text{pd}_A(M) \leq n$ by lemma 2.5.(2), therefore A is a strong n -perfect ring.
- 2) By Corollary 2.2 and 1) and this completes the proof of Theorem 2.3. ■

From this Theorem we deduce easily the following example of strong n -perfect rings.

Example 2.6 *Let D be an integral domain such that $\text{gldim}(D) = n$, let $K = \text{qf}(D)$ and let $n \geq 2$.*

Consider the quotient ring $S = K[X]/(X^n - X) = K + \overline{X}K[\overline{X}] = K + I$ with $I = \overline{X}K[\overline{X}]$. Set $R = D + I$. Then R is a strong n -perfect ring.

Proof. First we show that I is a pure ideal of R . Let α be an element of I , then $\alpha = \overline{X}^i(a_0 + a_1\overline{X} + \dots + a_{n-1}\overline{X}^{n-1})$ with $a_i \in K$ for $1 \leq i \leq n-1$, and $a_0 \neq 0$. As $\delta = a_0 + a_1\overline{X} + \dots + a_{n-1}\overline{X}^{n-1}$ is an invertible element.

$$\text{And } \overline{X}^i(1 - \overline{X}^{n-1}) = \overline{X}^i - \overline{X}^{n+(i-1)} = \overline{X}^i - \overline{X}^n\overline{X}^{i-1} = \overline{X}^i - \overline{X}^i = \overline{0}$$

Then $\alpha(1 - \overline{X}^{i-1}) = 0$ and $\overline{X}^{i-1} \in I$. Therefore I is a pure ideal of R by [12, Theorem 1.2.15].

On the other hand, we have S an artinian ring. According to [1, Corollary 28.8] we deduce S is a perfect ring and so S is a strong n -perfect ring, and also D is a strong n -perfect ring, since $\text{gldim}(D) = n$. Therefore R is a strong n -perfect ring by Theorem 2.3. ■

Later, we give an example showing that Theorem 2.3 is not true without assuming that the ideal Q is pure (see Example 2.11).

Now we link the strong n -perfect property with (n, d) -property in a particular cases. First, let n, d be positive integers, we say that R is an (n, d) -ring, if each n -presented R -module has projective dimension at most d , see for instance [4, 5, 7, 15]. In particular, the $(1, 0)$ -rings are the von Neumann regular rings, $(0, 1)$ -rings are the hereditary rings, $(1, 1)$ -rings are the semihereditary rings and $(0, 0)$ -rings are the semisimple rings [4, Theorem 1.3].

It is clear to see that if R is Noetherian, then R is an $(1, d)$ -ring if and only if R is an $(0, d)$ -ring. In the context of strongly d -perfect ring, we have:

Theorem 2.7 *Let R be a ring.*

R is an $(1, d)$ -ring and a strong d -perfect ring if and only if R is an $(0, d)$ -ring.

Proof. : Assume that R is an $(1, d)$ -ring and strong d -perfect. Hence $\text{wdim}(R) \leq d$ [12, Theorem 1.3.9] since for finitely generated I ideal of R , we have $\text{fd}(R/I) \leq d$ (since $\text{wdim}(R) \leq d$).

New let M be an R -module, thus $\text{fd}(M) \leq d$ (since $\text{wdim}(R) \leq d$) and so $\text{pd}(M) \leq d$ since R is a strong d -perfect ring. Therefore R is an $(0, d)$ -ring.

Conversely, assume that R is an $(0, d)$ -ring that is $\text{gldim}(R) \leq d$. It is clear that R is an $(1, d)$ -ring and a strong d -perfect ring. ■

According to proof of this theorem, we also show that strong n -perfect property links, in particular cases, the classical global and weak dimensions.

Corollary 2.8 *Let R be a strong n -perfect ring then: $\text{gldim}(R) \leq n$ if and only if $\text{wdim}(R) \leq n$.*

Since the $(1, 0)$ -rings are the von Neumann regular rings, the $(0, 0)$ -rings are the semisimple rings, the $(0, 1)$ -rings are the hereditary rings, and the $(1, 1)$ -rings are the semihereditary rings, then by Theorem 2.7 we have:

Corollary 2.9 1. *A von Neumann regular ring is a semisimple ring if and only if it is a perfect ring.*

2. *A semihereditary ring is an hereditary ring if and only if it is a strong 1-perfect ring.*

Remark 2.10 *There exists a ring R which is a von Neumann regular ring and an hereditary ring but is not a semisimple ring [4, Example 2.7]. By corollary 2.9, we deduce that R is a strong 1-perfect ring but it is not a perfect ring.*

Now we can show that Theorem 2.3 is not true without assuming that Q is a pure ideal.

Example 2.11 *Let $\mathbb{Z}$ denotes the ring of integers, and let $\mathbb{Q}$ denotes the field of rational numbers.*

Consider the discrete valuation domain $B = \mathbb{Q}[X]_{(X)} = \mathbb{Q} + XB$. Let $A = \mathbb{Z} + XB$, then.

1. B is a strong 1-perfect ring.
2. $A/XB \cong \mathbb{Z}$ is a strong 1-perfect ring.
3. XB is not a pure ideal, and A is not a strong 1-perfect ring.

Proof. : 1) We have B is a strong 1-perfect ring since B is an hereditary ring (corollary 2.9).

2) Similarly $A/XB \cong \mathbb{Z}$ is a strong 1-perfect ring.

3) We first set $M = XB$ and $E = A/M$ thus $A = \mathbb{Z} + M$ and $B = \mathbb{Q} + M$. The exact sequence of A -modules $0 \rightarrow M \rightarrow A \rightarrow \mathbb{Z} \rightarrow 0$ yields the exact sequence of A -modules

$$0 \rightarrow \text{Tor}_1^A(E, \mathbb{Z}) \rightarrow E \otimes_A M \rightarrow E \rightarrow E/ME \rightarrow 0$$

But $\text{Tor}_1^A(E, \mathbb{Z}) \cong E \otimes_A M \cong M/M^2 \neq 0$ Since M is principal and $ME = 0$, i.e., $\text{Tor}_1^A(A/M, \mathbb{Z}) \neq 0$. So M is not a pure ideal [12, Theorem 1.2.15].

On the other hand: we have $\text{fd}_A(E) \leq 1$, since $0 \rightarrow M \rightarrow A \rightarrow \mathbb{Z}(E) \rightarrow 0$ is exact and M is B -plat and B is A -plat.

If $\text{pd}_A(E) \leq 1$ was true, we would have an A -projective resolution:

$$0 \rightarrow P_1 \rightarrow P_0 \xrightarrow{u_0} E \rightarrow 0$$

Thus

$$(*) \quad 0 \rightarrow \text{Tor}_1^A(E, \mathbb{Z}) \rightarrow P_1 \otimes_A \mathbb{Z} \rightarrow P_0 \otimes_A \mathbb{Z} \xrightarrow{u_0 \otimes 1} E \otimes_A \mathbb{Z} \rightarrow 0$$

is exact. But $E \otimes_A \mathbb{Z} \cong E \otimes_A A/M \cong E/ME \cong E = \mathbb{Z}$, thus $E \otimes_A \mathbb{Z}$ is $\mathbb{Z}$ -projective, making $(*)$ split-exact and giving that $P_1 \otimes_A \mathbb{Z} \cong \text{Tor}_1^A(E, \mathbb{Z}) \oplus \text{Ker}(u_0 \otimes 1)$, so that $\text{Tor}_1^A(E, \mathbb{Z}) \cong M/M^2$ is a projective $\mathbb{Z}$ -module. But M/M^2 is a $\mathbb{Q}$ -vector space, so it would follow that $\mathbb{Q}$ is a projective $\mathbb{Z}$ -module, which is false since $\mathbb{Q} = qf(\mathbb{Z})$ and $\mathbb{Z} \neq \mathbb{Q}$. Thus $\text{pd}_A(E) > 1$. ■

But the transfer of the “ n -perfectness property” in pullback rings is not need to the hypothesis “ Q is a pure ideal” as follows to explain.

Proposition 2.12 *Let $A \hookrightarrow B$ be an injective flat ring homomorphism and let Q be an ideal of A such that $QB = Q$. Then:
If B is an n -perfect ring and A/Q is an n -perfect ring, then A is an n -perfect ring.*

Proof. Let E be a flat A -module then $E \otimes_A B$ is a flat B -module.

Thus $\text{pd}_B(E \otimes_A B) \leq n$ since B is an n -perfect ring.

On the other hand, $E \otimes_A A/Q$ is a flat A/Q -module, thus $\text{pd}_{A/Q}(E \otimes_A A/Q) \leq n$ since A/Q is an n -perfect ring. By lemma 2.4, we have $\text{pd}_R(E) \leq n$. Therefor A is an n -perfect ring. ■

In example 2.11, the ring A is an 1-perfect ring such that is not a strong 1-perfect ring. The following example shows that there exists the rings n -perfect are not a strong n -perfect rings for any n .

Example 2.13 *Let $\mathbb{C}$ denotes the field of complex numbers and $\mathbb{C}(X, Y)$ denotes the quotient field of the polynomial ring $\mathbb{C}[X, Y]$. Consider the discrete valuation domain $V = \mathbb{C}(X, Y)[Z]_{(Z)} = \mathbb{C}(X, Y) + ZV$.
Set the pullback ring $R = \mathbb{C}[X, Y] + ZV$. Then:*

1. R is an 2-perfect ring and it is not a strong 2-perfect ring .
2. $R[X_1]$ is an 3-perfect ring and it is not a strong 3-perfect.
3. Let $n \geq 4$ be integer $R[X_1, \dots, X_{n-2}]$ is an n -perfect ring and it is not strong n -perfect ring.

Proof. : By [6, Proposition 2.1] $\text{gldim}(R) = 3$ and $\text{wdim}(R) = 2$.

1) We have $R/ZV = \mathbb{C}[X, Y]$ is an 2-perfect ring (since $\text{gldim}(\mathbb{C}[X, Y]) = \text{wdim}(\mathbb{C}[X, Y]) = 2$) and V is an 2-perfect ring (since V is a discrete valuation domain). Hence, R is an 2-perfect ring (By proposition 2.12) and R is not a strong 2-perfect (by Corollary 2.8).

2) $R[X_1]$ is an 3-perfect ring (by [9, Example iv page 2]) and R is not strong 3-perfect (by $\text{wdim}(R[X_1]) = 3$ and $\text{gldim}(R[X_1]) = 4$).

3) By induction on n , we have for every $n \geq 4$, $R[X_1, \dots, X_{n-2}]$ is an n -perfect ring and it is not a strong n -perfect ring. ■

In the following we prove that the strong n -perfect property descends into a faithfully flat ring homomorphism.

Proposition 2.14 *Let $A \longrightarrow B$ be a ring homomorphism making B a faithfully flat A -module. if B is a strong n -perfect ring then A is a strong n -perfect ring.*

Proof. : Let M be an A -module such that $\text{fd}_A(M) \leq n$. Our aim is to show that $\text{pd}_A \leq n$.

Let $0 \longrightarrow P \longrightarrow F_{n-1} \longrightarrow \dots \longrightarrow F_1 \longrightarrow F_0 \longrightarrow M \longrightarrow 0$ be an exact sequence of A -modules, where F_i is a free A -module for each i , P is a flat A -module thus:

(*) $0 \longrightarrow P \otimes_A B \longrightarrow F_{n-1} \otimes_A B \longrightarrow \dots \longrightarrow F_1 \otimes_A B \longrightarrow F_0 \otimes_A B \longrightarrow M \otimes_A B \longrightarrow 0$ is an exact sequence of B -modules, where $F_i \otimes_A B$ is a free B -module for each i and $P \otimes_A B$ is a flat B -module, therefore $\text{fd}_B(M \otimes_A B) \leq n$ and so $\text{pd}_B(M \otimes_A B) \leq n$ B is a strong n -perfect. Hence, $P \otimes_A B$ is a projective B -module By (*).

According to [13, Example 3.1.4. page 82] we deduce that P is a projective A -module since P is a flat A -module. Then $\text{pd}_A(M) \leq n$ and so A is a strong n -perfect ring.

■

We profit by this proposition to study the strong n -perfect property of some particular rings.

Recall that the trivial extension of a ring R by an R -module M is the ring denotes by $R \ltimes M$ whose underling group is $R \times M$ with pairwise addition and multiplication given by $(a, e)(a', e') = (aa', ae' + a'e)$ (see for instance [11, 12, 14]).

Corollary 2.15 1. *Let A be a ring, let M be a flat A -module and let $R = A \ltimes M$ be the trivial ring extension of A by M . If R is a strong n -perfect ring then so is A .*

2. *Let $A \subset B$ be two ring such that B is a flat A -module. Let $S = A + XB[X]$, where X is an indeterminate over B . If S is a strong n -perfect ring then so is A .*

3. *Let R be a ring and X is an indeterminate over R . If $R[X]$ is a strong n -perfect ring then so is R .*

Proof. : 1) We have M a flat A -module, thus the ring homomorphism $A \hookrightarrow R$ is a faithfully flat, by Proposition 2.14, we conclude that A is a strong n -perfect ring since R is a strong n -perfect ring.

2) The ring B is a flat A -module and $XB[X] \cong B[X]$ thus $S = A + XB[X]$ is a faithfully flat A -module, also by Proposition 2.14 the ring A is a strong n -perfect since S is a strong n -perfect ring.

3) Similarly, by Proposition 2.14, since $R[X]$ is a faithfully flat R -module. ■

Next we establish the transfer of the strong n -perfect property to finite direct product.

Theorem 2.16 *Let $(R_i)_{i=1,\dots,m}$ be a family of rings. Then $\prod_{i=1}^m R_i$ is a strong n -perfect ring if and only if R_i is a strong n -perfect ring for each $i = 1, \dots, m$.*

The proof of the theorem involves the following results.

Lemma 2.17 [15, Lemma 2.5.(2)]

Let $(R_i)_{i=1,2}$ be a family of rings and E_i be an R_i -module for $i = 1, 2$. We have :
 $\text{pd}_{R_1 \times R_2}(E_1 \times E_2) = \sup\{\text{pd}_{R_1}(E_1), \text{pd}_{R_2}(E_2)\}.$

Lemma 2.18 *Let $(R_i)_{i=1,2}$ be a family of rings and E_i be an R_i -module for $i = 1, 2$. Then $\text{fd}_{R_1 \times R_2}(E_1 \times E_2) = \sup\{\text{fd}_{R_1}(E_1), \text{fd}_{R_2}(E_2)\}.$*

Proof. The proof is analogous to the proof of Lemma 2.17.

Proof of Theorem 2.16

By induction on m , it suffices to prove the assertion for $m = 2$. Let R_1 and R_2 be two rings such that $R_1 \times R_2$ is a strong n -perfect and let E_1 be an R_1 -module such that $\text{fd}_{R_1}(E_1) \leq n$, and let E_2 be an R_2 -module such that $\text{fd}_{R_2}(E_2) \leq n$. Then $\text{fd}_{R_1 \times R_2}(E_1 \times E_2) = \sup\{\text{fd}_{R_1}(E_1), \text{fd}_{R_2}(E_2)\}$ by lemma 2.18.

Thus $\text{pd}_{R_1 \times R_2}(E_1 \times E_2) \leq n$ since $R_1 \times R_2$ is a strong n -perfect ring.

But $\text{pd}_{R_1 \times R_2}(E_1 \times E_2) = \sup\{\text{pd}_{R_1}(E_1), \text{pd}_{R_2}(E_2)\}$ by lemma 2.17.

Therefore $\text{pd}_{R_1}(E_1) \leq n$ and $\text{pd}_{R_2}(E_2) \leq n$ and so R_1 and R_2 are a strong n -perfect rings .

Conversely, assume that R_1 and R_2 be two strong n -perfect rings. Let $E_1 \times E_2$ be an $R_1 \times R_2$ -module where E_i is an R_i -module for each $i = 1, 2$, such that $\text{fd}_{R_1 \times R_2}(E_1 \times E_2) \leq n$. Thus $\text{fd}_{R_1}(E_1) \leq n$ and $\text{fd}_{R_2}(E_2) \leq n$ (by lemma 2.18) and so $\text{pd}_{R_1}(E_1) \leq n$ and $\text{pd}_{R_2}(E_2) \leq n$ (since R_1 and R_2 are strong n -perfect rings). Therefore $\text{pd}_{R_1 \times R_2}(E_1 \times E_2) \leq n$ (By Lemma 2.17) and so $R_1 \times R_2$ is a strong n -perfect rings. ■

Recall that a ring R is called an S -ring if every finitely generated flat R -module is projective (see [17]).

The object of this remark is to conclude, in general case, that the strong n -perfect property is not characterized by the cyclic modules.

Remark 2.19 *The ring A of Example 2.11 is an integral domain, hence, by [8, page 288], the ring A is an S -ring, and by [19, Corollary 1.7] we deduce on the ring A every cyclic flat module is projective. But according also to Example 2.11 the ring A is not perfect since it is not strong 1-perfect. Therefore the perfect property is not characterized by the cyclic modules.*

References

- [1] F. W. Anderson. K. R. Fuller; *Rings and Categories of Modules*, Springer-Verlag, 2nd edition, in: Graduate Texts Mathematics, vol. 13, New York, (1992).
- [2] H. Bass; *finitistic dimension and a homological generalization of semi-primary rings*, Trans. Amer. Math. Soc. 95 (1960), 466-488.
- [3] H. Cartan and S. Eilenberg; *Homological algebra*, Princeton University press, 1956.
- [4] D. L. Costa; *Parameterizing families of non-noetherian rings*, Comm. Algebra 22 (1994), 3997-4011.
- [5] D. L. Costa S. D. Kabbaj; *Classes of $D + M$ ring defined by homolgical conditions*, Comm. Algebra 24 (1996), 891-906.
- [6] D. E. Dobbs; *On the global dimensions of $D + M$* , Canad. Math. Bull. 18 (1975), 657-660.
- [7] D. E. Dobbs, S. Kabbaj, N. Mahdou, and M. Sobrani; *When is $D + M$ n -coherent and an (n, d) -domain?*, Lecture Notes in Pure and Appl. Math., Dekker, 250 (1999), 257-270.
- [8] S. Endo, *On flat modules over commutative rings*, J. Math. Soc. Japan 14 (1962), 284-291.
- [9] E. E. Enochs, O. M. G. Jenda, and J. A. López-Romos, *Dualizing modules and n -perfect rings*, Proc. Edinb. Math. Soc. (2) 48 (2005), no. 1, 75-90.
- [10] M. Fontana, *Topologically defined classes of commutative rings*, Ann. Mat. Pura Appl. 123 (1980), 331-355.
- [11] R. M. Fossum, P. A. Griffith, and I. Riten; *Trivial extensions of abelian categories*, Springer-Verlag, Lecture Notes in Mathematics, (1975) 456.

- [12] S. Glaz; *Commutative Coherent Rings*, Springer-Verlag, Lecture Notes in Mathematics, 1371 (1989).
- [13] L. Gruson and M Raynaud, *Critères de platitude et de projectivité. Techniques de platification d'un module*, Inv. Math. 13 (1971), 1-89.
- [14] Huckaba, J, A. *Commutative Coherent Rings with Zero Divisors*, Marcel Dekker, New York Basel, 1989.
- [15] N. Mahdou; *On Costa's conjecture*, Comm. Algebra 29 (7) (2001), 2775-2785.
- [16] Osborne, M. S. *Basic Homological Algebra*, Springer-Verlag, Graduate Text in Mathematics, New York, Berlin, 196. (2000).
- [17] G. Puninski, P. Rothmaler, *When every finitely generated flat module is projective* , J. Algebra 277 (2004) 542-558.
- [18] J. Rotman; *An introduction to homological algebra*, Press, New York, 25(1979).
- [19] W.V. Vasconcelos, *On finitely generated flat module*, Tran. Amer. Math. Soc. 138 (1969) 505-512.